

Taking VYZULTA as prescribed is important for your glaucoma treatment

VYZULTA.
(latanoprostene
bunod ophthalmic
solution), 0.024%



WHAT is glaucoma?

Glaucoma is a **persistent eye disease** in which high eye pressure can lead to permanent vision loss if not treated



HOW can VYZULTA help?

VYZULTA can **help lower your eye pressure** by increasing drainage through two different pathways in the eye



WHY is lowering eye pressure important?

Lowering eye pressure is an important step in **treating glaucoma and helping to preserve eyesight**



You can make a difference
Use your drops every day as prescribed

INDICATION

VYZULTA® (latanoprostene bunod ophthalmic solution), 0.024%, is used to lower intraocular (eye) pressure in patients with open-angle glaucoma or ocular hypertension.

IMPORTANT SAFETY INFORMATION

- Brown color, or increased brown color, of the iris can occur, which may be permanent. Darkening of the eyelid skin may also occur, which is usually reversible after VYZULTA treatment is discontinued.
- Gradual changes to eyelashes may occur, including increased length, thickness, color, number of eyelashes and/or direction of eyelash growth. These changes may result in a difference in appearance of each eye. These changes are usually reversible after VYZULTA treatment is discontinued.
- Avoid touching the tip of the container to the eye or any other surface, including a finger, in order to avoid contamination of the solution by bacteria. Serious damage to the eye and possible loss of vision may result from using contaminated solutions.

Please see additional Important Safety Information on the reverse side and accompanying full Prescribing Information in pocket.

Continue for helpful tips on taking VYZULTA every day

3 steps to help you remember to take VYZULTA every day


VYZULTA.
(latanoprostene buno-d ophthalmic solution), 0.024%



PLAN

Keep drops in the same spot and take them at the same time every night



REMIND

Use physical notes at home or set reminders with your mobile or smart home devices



REPEAT

Make taking VYZULTA part of other routine evening activities like brushing your teeth or washing your face



REMEMBER: Follow up with your doctor and obtain refills when necessary. Ask your doctor if a coupon may be available to you

IMPORTANT SAFETY INFORMATION (Cont.)

- Tell your doctor immediately about any change in your eye, any eye condition that develops, any sudden change in your vision, any eye side effects, or if you have eye surgery.
- Remove contact lenses before putting VYZULTA in your eyes. Lenses may be reinserted 15 minutes later.
- If you are using other eye drops in addition to VYZULTA, wait 5 minutes between using each type of eye drop.
- The most common side effects seen in studies were eye redness, eye irritation, eye pain, and pain after putting the drops in your eye.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see accompanying full Prescribing Information in pocket.

BAUSCH+LOMB

VYZULTA and the V design are trademarks of Bausch & Lomb Incorporated or its affiliates. ©2021 Bausch & Lomb Incorporated or its affiliates. All rights reserved. VYZ.0014.USA.21